

Initial conditions for hydrodynamics: implications for phenomenology*

WOJCIECH BRONIEWSKI^{a,b}, WOJCIECH FLORKOWSKI^{a,b},
MIKOŁAJ CHOJNACKI^a, AND ADAM KISIEL^{c,d}

^aThe H. Niewodniczański Institute of Nuclear Physics,
Polish Academy of Sciences, PL-31342 Kraków, Poland

^bInstitute of Physics, Jan Kochanowski University, PL-25406 Kielce, Poland

^cFaculty of Physics, Warsaw University of Technology, PL-00661 Warsaw, Poland

^dDepartment of Physics, Ohio State University, Columbus, OH 43210, USA

(Received 29 December 2008)

It is shown how the initial azimuthally asymmetric flow develops from the free-streaming + sudden equilibration approximation to the early dynamics in relativistic heavy-ion collisions. Consequences for the hydrodynamics and physical results are discussed.

PACS numbers: 25.75.-q, 25.75.Dw, 25.75.Ld

We present a description of the early-stage dynamics in relativistic heavy ion collisions, where the free-streaming (FS) of partons is followed by a sudden equilibration (SE) to a thermalized phase, providing initial conditions for the subsequent hydrodynamic evolution. This FS+SE approximation has been proposed by Kolb, Sollfrank, and Heinz [1]. It has been further discussed in the context of the isotropization problem by Jas and Mrówczyński [2], as well as used to analyze the early development of collective flow by Sinyukov, Gyulassy, Karpenko, and Nazarenko [3, 4, 5]. In this talk we point out the emergence of the *initial azimuthally asymmetric* flow from FS+SE.

The cartoon of the approach is given in Fig. 1. Rather than assuming a gradual transition from an inequilibrium partonic stage to a thermalized system (top panel), in FS+SE one approximates this early stage of evolution with free streaming followed by a sudden equilibration (bottom panel). Note that this is analogous to the standard treatment of the freeze-out at the end of the hydrodynamic evolution, where the continuous decoupling of

* Presented by WB at the IV Workshop on Particle Correlations and Femtoscopy, Cracow, 11-14 September 2008




Fig. 1. Evolution of the system formed in relativistic heavy-ion collisions, consisting of partonic free streaming, equilibration, hydrodynamics, freeze-out, and free streaming of hadrons to detectors. Top: equilibration and freeze-out occur gradually. Bottom: the approximation of the sudden equilibration and instantaneous freeze-out.

hadrons (see the talks by Yu. Sinyukov and J. Knoll in these proceedings) is approximated with a sudden Cooper-Frye freeze-out.

Ever since the FS+SE has been proposed, it has been generally thought that it unavoidably reduces the elliptic flow, which develops hydrodynamically due to the azimuthal asymmetry of the initial density profile for non-central collisions. Admittedly, FS decreases the spatial azimuthal asymmetry of the system with time. We carefully reexamine this argument. The point is that while FS *alone* obviously cannot generate azimuthal asymmetry in the momentum distribution, due to the well-known fact that interactions among produced particles are needed to generate v_2 , the SE in fact does the job. This is because SE is dynamical in nature, resulting in an abrupt change, due to interactions, of the energy-momentum tensor of the system into a diagonal form (in the reference frame of the fluid element) of perfect hydro.

One may interpret the FS+SE approach as an approximation to viscous hydrodynamics. Instead of considering a complicated viscous system far from the thermal equilibrium, where the partonic cross section has a finite value, one initially treats the partons as free, and later supplies them with a large cross section which results in an instantaneous equilibration of the system and transition to perfect hydrodynamics. Such an interpretation works when viscosity decreases with time, or equivalently, the partonic cross section increases. Confinement effects, which switch on as the distance between the partons increases, provide a mechanism for this behavior and support this interpretation, although a convincing solution of the early thermalization or isotropization problems is still missing despite many theoretical efforts.

Here we analyze a boost-invariant system with an initial Gaussian transverse energy profile, $n(x_0, y_0) = \exp\left(-\frac{x_0^2}{2a^2} - \frac{y_0^2}{2b^2}\right)$, where the widths a and

b depend on centrality and are obtained with GLISSANDO [6]. We introduce the initial, $\tau_0 = \sqrt{t_0^2 - z_0^2}$, and final, $\tau = \sqrt{t^2 - z^2}$, proper times of free streaming, as well as the space-time rapidities $\eta_0 = \frac{1}{2} \log \frac{t_0 - z_0}{t_0 + z_0}$ and $\eta = \frac{1}{2} \log \frac{t - z}{t + z}$. Elementary kinematics, following from the fact that a massless parton moves along a straight line with the velocity of light and a four momentum $p^\mu = (p_T \cosh Y, p_T \cos \phi, p_T \sin \phi, p_T \sinh Y)$, relates the initial and final coordinates of the parton:

$$\begin{aligned} \tau \sinh(\eta - Y) &= \tau_0 \sinh(\eta_0 - Y), \quad x = x_0 + d \cos \phi, \quad y = y_0 + d \sin \phi, \\ d &= \frac{t - t_0}{\cosh Y} = \tau \cosh(Y - \eta) - \sqrt{\tau_0^2 + \tau^2 \sinh^2(Y - \eta)}. \end{aligned} \quad (1)$$

Consequently, the phase-space densities of partons at the proper times τ_0 and τ are related,

$$\begin{aligned} \frac{d^6 N(\tau)}{dY d^2 p_T d\eta dx dy} &= \int d\eta_0 dx_0 dy_0 \frac{d^6 N(\tau_0)}{dY d^2 p_T d\eta_0 dx_0 dy_0} \times \\ &\delta(\eta_0 - Y - \operatorname{arcsinh}[\frac{\tau}{\tau_0} \sinh(\eta - Y)]) \delta(x - x_0 - d \cos \phi) \delta(y - y_0 - d \sin \phi). \end{aligned} \quad (2)$$

We assume for simplicity a factorized form of the initial parton distribution,

$$\frac{d^6 N(\tau_0)}{dY d^2 p_T d\eta_0 dx_0 dy_0} = n(x_0, y_0) F(Y - \eta_0, p_T). \quad (3)$$

In Ref. [7] we show that to a very good accuracy $F(Y - \eta_0, p_T) \sim \delta(Y - \eta)$. Then, the energy-momentum tensor of the system at $\eta = 0$ is

$$\begin{aligned} T^{\mu\nu} &= A \int_0^{2\pi} d\phi n(x - (\tau - \tau_0) \cos \phi, y - (\tau - \tau_0) \sin \phi) \times \\ &\begin{pmatrix} 1 & \cos \phi & \sin \phi & 0 \\ \cos \phi & \cos^2 \phi & \cos \phi \sin \phi & 0 \\ \sin \phi & \cos \phi \sin \phi & \sin^2 \phi & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \end{aligned} \quad (4)$$

where A is a constant from the p_T integration. Next, at each point we pass to the local reference frame in which the T_{0i} components of the energy-momentum tensor vanish. The four-velocity needed for the appropriate boost is found from the condition $T^{\mu\nu}(x, y) u_\nu(x, y) = \varepsilon(x, y) g^{\mu\nu} u_\nu(x, y)$, where ε is the energy density in the local rest frame. In the left part of Fig. 2 we show the profile of ε together with the energy-momentum tensor in the local rest frame (in units of ε), displayed at a few points. We note that it has the structure very close to the case of the *perfect transverse hydrodynamics* of massless particles [8], where the transverse pressure is equal




Fig. 2. Left: Energy-density contours with the energy-momentum tensor in the local rest frame (in units of ε) shown at a few points (dots). Right: the profile of the transverse velocity, $v = \sqrt{v_x^2 + v_y^2}$, in units of c .




Fig. 3. Left: The schematic development of the partonic elliptic flow ϵ_p from FS+SE. Right: the value of the generated momentum asymmetry ϵ_p plotted as a function of the proper time when the Landau matching is imposed. The dotted (dashed) line corresponds to matching to isotropic (transverse) hydro, while the solid line shows the results of hydrodynamics only, with no FS present. The top curve shows the spatial asymmetry ϵ , which decreases with time.

to $\varepsilon/2$. Small departures from this form, present in our case, have the structure of the shear tensor used to include the viscosity effects in transverse hydrodynamics [9].

The right part of Fig. 2 shows the transverse velocity profile. We note that it is azimuthally asymmetric (stronger in the direction of the impact parameter), which simply reflects the original geometry. In fact, for low free-streaming times and close to the origin one finds $\mathbf{v} = -\frac{1}{3}(\tau - \tau_0)\nabla n/n$.




Fig. 4. The pion HBT radii R_{side} , R_{out} , R_{long} , and the ratio $R_{\text{out}}/R_{\text{side}}$ for central collisions. The darker (lighter) lines describe the results with (without) FS+SE. The data from Ref. [10].

Thus the space – collective velocity correlations are induced.

Following Ref. [1], we now consider a convenient measure of the *momentum* anisotropy, $\epsilon_p = (\langle T_{xx} \rangle - \langle T_{yy} \rangle) / (\langle T_{xx} \rangle + \langle T_{yy} \rangle)$, where brackets denote the spatial integration. In the FS phase identically $\epsilon_p = 0$, as no interactions have occurred. Then, at the proper time τ , SE occurs. As a result, the energy momentum tensor in the local frame is replaced, due to interactions, with a diagonal form of the perfect hydrodynamics: $T^{\mu\nu} \rightarrow \text{diag}(\epsilon, \epsilon/3, \epsilon/3, \epsilon/3)$ for the isotropic, or $T^{\mu\nu} \rightarrow \text{diag}(\epsilon, \epsilon/2, \epsilon/2, 0)$ for the transverse hydro. It is this *Landau matching* condition which causes ϵ_p to *jump to a nonzero value*. We show the result in Fig. 3.

Finally, we compare the physical results obtained with (isotropic) hydrodynamics only (starting at an early proper time $\tau_0 = 0.25$ fm) to the results obtained with FS from τ_0 to $\tau = 1.0$ fm, followed by SE and (isotropic) hydro. We use the realistic equation of state [11]. The statistical hadronization is carried out with the help of THERMINATOR [12]. In Fig. 4 we notice strikingly similar results for the two considered cases, not to mention the very good description of the HBT data, including the infamous ratio $R_{\text{out}}/R_{\text{side}}$. The similarity and agreement is similar for the p_T -spectra and v_2 [7, 11], as well as for other centrality classes. In Ref. [13] we have also shown that our model calculations reproduce very well the azHBT pion interferometry.

A practical conclusion from our study is that the inclusion of FS+SE may be used to *delay the start of perfect hydrodynamics* to “comfortable” times of about 1 fm/c. The physical results remain practically unaltered, since the decrease of the spatial anisotropy with time, resulting in milder hydrodynamic development of v_2 , is intertwined with the buildup of the *initial azimuthally asymmetric* flow. Recall that in some studies, in order to obtain a proper description of the particle spectra and femtoscopy, hydro was used with initialization times of 0.1 fm/c [14, 15] (with viscous hydro). Recently, the phenomenological relevance of the initial flow, first examined in hydrodynamics in Ref. [16], has been emphasized in Refs. [15, 17] (see also the contribution of S. Pratt to these proceedings).

Two of us (WB and WF) are grateful to Piotr Bożek and Stanisław Mrówczyński for useful conversations.

REFERENCES

- [1] P.F. Kolb, J. Sollfrank and U.W. Heinz, Phys. Rev. C62 (2000) 054909, hep-ph/0006129.
- [2] W. Jas and S. Mrowczynski, Phys. Rev. C76 (2007) 044905, 0706.2273.
- [3] Y.M. Sinyukov, Acta Phys. Polon. B37 (2006) 3343.
- [4] M. Gyulassy et al., Braz. J. Phys. 37 (2007) 1031.
- [5] Y. Sinyukov, (2008), talk presented at Quark Matter 2008, Jaipur, India, 4-10 February 2008.
- [6] W. Broniowski, M. Rybczynski and P. Bozek, Comput. Phys. Commun. 180 (2009) 69, 0710.5731.
- [7] W. Broniowski et al., (2008), 0812.3393.
- [8] A. Bialas, M. Chojnacki and W. Florkowski, Phys. Lett. B661 (2008) 325, 0708.1076.
- [9] P. Bozek, Acta Phys. Polon. B39 (2008) 1375, 0711.2889.
- [10] STAR, J. Adams et al., Phys. Rev. C71 (2005) 044906, nucl-ex/0411036.
- [11] W. Broniowski et al., Phys. Rev. Lett. 101 (2008) 022301, 0801.4361.
- [12] A. Kisiel et al., Comput. Phys. Commun. 174 (2006) 669, nucl-th/0504047.
- [13] A. Kisiel et al., (2008), 0808.3363.
- [14] S. Pratt and J. Vredevoogd, Phys. Rev. C78 (2008) 054906, 0809.0516.
- [15] S. Pratt, (2008), 0811.3363.
- [16] M. Chojnacki, W. Florkowski and T. Csorgo, Phys. Rev. C71 (2005) 044902, nucl-th/0410036.
- [17] M.A. Lisa and S. Pratt, (2008), 0811.1352.